



Product Information: Recombinant human Activin A, 10µg

1. Introduction

Activin A is a member of the TGF-beta superfamily of cytokines and is involved in a wide range of biological processes including tissue morphogenesis and repair, fibrosis, inflammation, neural development, hematopoiesis, reproductive system function, and carcinogenesis. Human Activin A is a 26.0 kDa disulfide-linked homodimer of two βA chains, each containing 116 amino acid residues. Activin A is mainly used for stem cell cultivation in order to differentiate the stem cell into endoderm or mesoderm.

Recombinate Human Activin A

Let ACTORODIE

Let ACT

Source: *Escherichia coli*Amino Acids Sequence:

GLECDGKVNI CCKKQFFVSF KDIGWNDWII APSGYHANYC EGECPSHIAG TSGSSLSFHS
TVINHYRMRG HSPFANLKSC CVPTKLRPMS MLYYDDGQNI IKKDIQNMIV EECGCS

2. Materials Provided

Component	Volume	Storage
Recombinant human Activin A, 10µg (0.1 mg/ml)	100 µl	Store at -80°C

3. Preparation for Use

Recombinant human Activin A is provided frozen and can be stored at -80°C until use. Before use, this frozen product should be thawed on ice. Avoid repeated freeze-thaw cycles. As a general guide, if product is dispensed after the first thawing, do not store the aliquots at -20°C for more than 6 months (within 4 years from the manufacturing date) or at -80°C for more than 4 years from the manufacturing date.

4. Precaution and Disclaimer

Recombinant human Activin A is for research use only and is not for diagnostic use, therapeutic use (including clinical research), or as a food. Activin A is dissolved in 50 mM sodium acetate buffer, pH4.6 (0.2 µm filtered).

5. Contact the following department for product information:

Business Strategy & Planning Dept., AminoScience Division, AJINOMOTO CO., INC.

1-15-1 Kyobashi, Chuo-ku, Tokyo 104-8315, Japan

E-mail: stemfit@ajinomoto.com

For Research Use Only. Not Intended for Human or Animal Diagnostic or Therapeutic Uses

Version 3 (Nov 2020)



Email: info@nacalaiusa.com

TEL: (858) 404 0403 www.nacalaiusa.com